

University of Groningen

Ultrafast ignition of a uni-directional molecular motor

Conyard, Jamie; Addison, Kiri; Heisler, Ismael A.; Cnossen, Arjen; Browne, Wesley R.; Meech, Stephen R.; Feringa, Bernard

Published in:

Proceedings of the XVIIIth International Conference on Ultrafast Phenomena

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2013

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Conyard, J., Addison, K., Heisler, I. A., Cnossen, A., Browne, W. R., Meech, S. R., & Feringa, B. (2013). Ultrafast ignition of a uni-directional molecular motor. In M. Chergui, S. Cundiff, A. Taylor, R. D. Vivie-Riedle, & K. Yamanouchi (Eds.), *Proceedings of the XVIIIth International Conference on Ultrafast Phenomena* (41 ed., Vol. 41, pp. 05016-p.1-05016-p.3). University of Groningen, Stratingh Institute for Chemistry.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Ultrafast ignition of a uni-directional molecular motor

Jamie Conyard,¹ Kiri Addison,¹ Ismael A. Heisler,¹ Arjen Cnossen,² Wesley R. Browne,² Ben L. Feringa² and Stephen R. Meech¹


^aSchool of Chemistry, University of East Anglia, Norwich Research Park, Norwich NR4 7TJ

^bStratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747AG Groningen, The Netherlands

Abstract. Light-driven molecular motors convert light into mechanical energy via excited state reactions. In this work we follow sub-picosecond primary events in the cycle of a two-stroke unidirectional motor by fluorescence up-conversion and transient absorption.

1 Introduction

Light driven molecular machines are designed to convert photon energy into mechanical energy, which can often be achieved through an excited state isomerisation reaction. For the operation of molecular motors it is also clearly essential to exercise control over the direction of motion. Unidirectional operation of light-driven molecular motors was first reported by Feringa and co-workers in their studies of chiral overcrowded alkenes (Fig. 1)[1]



2 Results and Discussion

Time resolved fluorescence was measured at wavelengths spanning the emission spectrum. The decay on blue edge is ultrafast, revealing components on the order of 100 fs. As the observation wavelength increases the decay time lengthens, but no risetime is registered on the red edge of the spectrum (Fig. 2). Superimposed on the fast fluorescence decay is an oscillatory component which can be analysed in terms of one highly damped (90 fs) low frequency mode (113 cm^{-1}) and one less damped (220 fs) higher frequency mode (180 cm^{-1}); these are illustrated in the Fourier transform (inset). Such oscillations in the fluorescence are characteristic of vibrational modes in the excited state which are displaced with respect to the ground state. Ultrafast fluorescence is particularly well suited to the detection of such excited state modes




Figure 2. Time resolved fluorescence observed for the motor molecule of Fig. 1. Wavelengths span the spectrum from 455 nm (dark blue) to 556 nm (red). The inset shows a Fourier transform of the oscillatory part of the fluorescence.

The wavelength resolved time resolved fluorescence can be converted into time dependent emission spectra, which reveal that the spectral mean frequency shifts to the red as the integrated emission intensity collapses on a time scale of a few hundred femtoseconds. Subsequent to these fast events the fluorescence decays with a time constant of 1.4 ps. The oscillatory features evident in the wavelength resolved data of Fig. 2 translate to an oscillation in both the intensity and the mean frequency of the time resolved spectrum[2]. This shows that the low frequency modes excited are coupled to the electronic transition moment.

Transient absorption measurements reveal two spectral features (Fig. 3) with distinct kinetics. The lowest energy one is formed rapidly (within the 300 fs time resolution of the apparatus) and




Figure 3. Transient absorption data for the motor in the visible region

decays with a 1.5 ± 0.3 ps decay constant. The higher energy feature is evidently a mixture of the 1.5 ps transient, stimulated emission a very long lived (> 100 ps) transient and a ground state bleach. We assign the 1.5 ps state to the relaxed excited state observed in the fluorescence data.

Based on these data we assign the primary events in the photoactivated stroke of a molecular motor following electronic excitation as follows. The bright Franck-Condon excited state relaxes in a few hundred femtoseconds to a less bright relaxed state. The high rate suggests a small scale energetically downhill structural transformation, possible involving pyramidalization at the C9 carbon, as suggested by quantum chemical calculation[3]. The relaxed state populated in ca 200 fs and relaxes back to the ground state in 1.5 ps. Coherently excited vibrational modes modulate the transition moment, connecting the dark and bright (F-C) states. These assignments are illustrated schematically in Fig. 4. It is interesting that the coherently excited low frequency modes relax on the sub 100 fs time scale, setting an upper limit to the time available for exerting coherent control over the primary photochemical step.




Figure 4. Schematic representation of the primary steps in a molecular motor

One route to the optical control of motor efficiency is through shaped light pulses, which may direct the excited state population towards or away from a conical intersection. However, the very fast damping of the lowest frequency modes suggests that the timescale in which this can be achieved is small. As an alternative to optical control of the rate and efficiency of motor operation we have investigated the effect of chemical modulation. Electron withdrawing (cyano, chloro) and donating (methoxy) groups were substituted on the 4' position on the naphthalenic ring. The effect of the substitution on the ground state structure was assessed. The data are complex but electron withdrawing groups accelerate the decay of the FC state, but extend the lifetime of the dark state. These data are being analysed in terms of modifications to the shape of the PES such as that described in Figure 4.

References

- [1] N. Koumura, R. W. J. Zijlstra, R. A. van Delden, N. Harada, and B.L. Feringa Light-driven monodirectional molecular rotor. *Nature* **401** (6749), 152 (1999)
- [2] J. Conyard, K. Addison, I. A. Heisler, A. Cnossen, W. R. Browne, B.L. Feringa and S. R. Meech Ultrafast Dynamics in the Ignition Phase of Molecular Rotary Motors, *Nature Chem.* **4**, 457 (2012).
- [3] Kazaryan, A. *et al.*, Understanding the Dynamics Behind the Photoisomerization of a Light-Driven Fluorene Molecular Rotary Motor. *J. Phys. Chem. A* **114**, 5058 (2010)